

Entanglement purification through Zeno-like measurements

K. YUASA, H. NAKAZATO and M. UNOKI

Department of Physics, Waseda University, Tokyo 169-8555, Japan

E-mail: yuasa@hep.phys.waseda.ac.jp

Abstract. We present a novel method to purify quantum states, i.e. *purification through Zeno-like measurements*, and show an application to *entanglement purification*.

PACS numbers: 03.65.Xp, 03.67.Mn

1. Introduction

One of the main obstacles to the experimental realisation of the ideas of quantum computation and quantum information is ‘decoherence’ [1, 2]. There are many attempts to overcome this problem, and several approaches are proposed for it. Among them are the ‘purification/distillation’ technologies [2, 3], i.e. methods to extract pure states, especially entanglements, from general mixed states recovering/preparing quantum coherence. Contributing to this subject, we have recently proposed a novel method to purify quantum states, i.e. purification through Zeno-like measurements [4]. Here we show that it is possible to apply it for extracting *entanglement*, which is the key resource to the fundamentals of quantum computation and information.

2. Purification through Zeno-like measurements

Let us recapitulate the framework of our purification [4]. We consider two quantum systems X and A interacting with each other. The total system is initially in a general *mixed* state ϱ_{tot} , from which we try to extract a pure state in A. We first make a measurement on X *to confirm that it is in a given state* $|\phi\rangle_X$. If it is found in the state $|\phi\rangle_X$, the state of the total system is projected by the projection operator

$$\mathcal{O} = |\phi\rangle_X \langle\phi| \otimes \mathbb{1}_A. \quad (1)$$

We then let the total system start to evolve under the Hamiltonian H_{tot} and repeat the same measurement on X at time intervals τ . After N repetitions of successful confirmations, the state of the total system, $\varrho_{\text{tot}}^{(\tau)}(N)$, is given by

$$\varrho_{\text{tot}}^{(\tau)}(N) = (\mathcal{O} e^{-iH_{\text{tot}}\tau} \mathcal{O})^N \varrho_{\text{tot}} (\mathcal{O} e^{iH_{\text{tot}}\tau} \mathcal{O})^N / P^{(\tau)}(N) = |\phi\rangle_X \langle\phi| \otimes \varrho_A^{(\tau)}(N), \quad (2)$$

$$\varrho_A^{(\tau)}(N) = (V_\phi(\tau))^N \varrho_A (V_\phi^\dagger(\tau))^N / P^{(\tau)}(N), \quad (3)$$

where $\varrho_A = {}_X \langle\phi| \varrho_{\text{tot}} |\phi\rangle_X$ is the state of A after the zeroth confirmation,

$$V_\phi(\tau) \equiv {}_X \langle\phi| e^{-iH_{\text{tot}}\tau} |\phi\rangle_X \quad (4)$$

is a projected time-evolution operator, and $P^{(\tau)}(N)$ is the normalisation factor,

$$P^{(\tau)}(N) = \text{Tr}[(\mathcal{O}e^{-iH_{\text{tot}}\tau}\mathcal{O})^N \varrho_{\text{tot}}(\mathcal{O}e^{iH_{\text{tot}}\tau}\mathcal{O})^N] = \text{Tr}_A[(V_\phi(\tau))^N \varrho_A(V_\phi^\dagger(\tau))^N]. \quad (5)$$

Note that we retain only the events where X is found in the state $|\phi\rangle_X$ at *every* measurement; other events, resulting in failure to purify A, are discarded. The normalisation factor $P^{(\tau)}(N)$ in (5) is nothing but the probability for the *successful events* and is the probability of obtaining the state given in (2) and (3).

Let us assume for simplicity that the eigenvalues λ_n of $V_\phi(\tau)$, which are complex-valued in general, are discrete and $V_\phi(\tau)$ is decomposed (diagonalised) as

$$V_\phi(\tau) = \sum_n \lambda_n |u_n\rangle_A \langle v_n|, \quad (6)$$

where $|u_n\rangle_A$ and ${}_A\langle v_n|$ are the right and left eigenvectors of $V_\phi(\tau)$, respectively, belonging to the eigenvalue λ_n and satisfy ${}_A\langle v_n|u_m\rangle_A = \delta_{mn}$. (The right eigenvectors are normalised as ${}_A\langle u_n|u_n\rangle_A = 1$ in the following.) It is now easy to observe the asymptotic behaviour of the state of A in (3). Since the eigenvalues λ_n are bounded as $0 \leq |\lambda_n| \leq 1$, each term in the expansion of $(V_\phi(\tau))^N$ decays away and a single term dominates asymptotically as the number of measurements, N , increases,

$$(V_\phi(\tau))^N = \sum_n \lambda_n^N |u_n\rangle_A \langle v_n| \rightarrow \lambda_0^N |u_0\rangle_A \langle v_0| \quad \text{as } N \text{ increases}, \quad (7)$$

provided the largest (in magnitude) eigenvalue λ_0 is unique, discrete and non-degenerate. Then, the state of A in (3) accordingly approaches the pure state $|u_0\rangle_A$,

$$\varrho_A^{(\tau)}(N) \rightarrow |u_0\rangle_A \langle u_0| \quad \text{as } N \text{ increases}. \quad (8)$$

This is the purification we have found recently [4]: extraction of a pure state $|u_0\rangle_A$ from an *arbitrary* mixed state ϱ_A through repeated measurements on X. Since we repeat measurements (on X) as in the case of the quantum Zeno effect [5], we call such measurements ‘Zeno-like measurements’.[‡]

The above observation shows that the assumption of the ‘spectral decomposition’ (6) is not essential but the existence of the *unique, discrete and nondegenerate largest eigenvalue* λ_0 is crucial for the purification. Furthermore, note the asymptotic behaviour of the ‘success probability’ $P^{(\tau)}(N)$: it decays asymptotically as

$$P^{(\tau)}(N) \rightarrow |\lambda_0|^{2N} {}_A\langle v_0|\varrho_A|v_0\rangle_A \quad \text{as } N \text{ increases}, \quad (9)$$

which is dominated by the eigenvalue λ_0 . Efficient purification is possible if it satisfies the condition $|\lambda_0| = 1$, which suppresses the decay in (9). At the same time, if the other eigenvalues are much smaller than λ_0 in magnitude, purification is achieved faster. Hence,

$$|\lambda_0| = 1, \quad |\lambda_n/\lambda_0| \ll 1 \quad \text{for } n \neq 0 \quad (10)$$

are the conditions for the *efficient purification*, which we try to achieve by adjusting parameters such as τ , $|\phi\rangle_X$ and those in the Hamiltonian H_{tot} .

[‡] It should be noted however that the time interval τ in this scheme is not necessarily small as in the ordinary Zeno measurements, and the purification (8) is not due to the quantum Zeno effect.

3. Application to entanglement purification

Now we show an interesting application of the above scheme: application to *entanglement purification*. Let us consider, for example, a three-qubit system with the Hamiltonian

$$H_{\text{tot}} = \Omega \frac{1 + \sigma_3^X}{2} + \Omega \frac{1 + \sigma_3^A}{2} + \Omega \frac{1 + \sigma_3^B}{2} + g(\sigma_+^X \sigma_-^A + \sigma_-^X \sigma_+^A) + g(\sigma_+^X \sigma_-^B + \sigma_-^X \sigma_+^B), \quad (11)$$

where σ_i ($i = 1, 2, 3$) are the Pauli operators and $\sigma_{\pm} = (\sigma_1 \pm i\sigma_2)/2$ are the ladder operators. Qubit X is coupled to qubits A and B. We confirm X to be in the state $|\phi\rangle_X$ repeatedly at time intervals τ and end up with an extraction of an entanglement between A and B, which are initially in a general mixed state ϱ_{tot} .

The projected time-evolution operator $V_{\phi}(\tau)$ is given, in this case, by

$$\begin{aligned} V_{\phi}(\tau) = & |\Psi^-\rangle_{AB} \langle \Psi^-| (|\beta|^2 + |\alpha|^2 e^{-i\Omega\tau}) e^{-i\Omega\tau} \\ & + |\downarrow\downarrow\rangle_{AB} \langle \downarrow\downarrow| (|\beta|^2 + |\alpha|^2 e^{-i\Omega\tau} \cos \sqrt{2}g\tau) \\ & + |\Psi^+\rangle_{AB} \langle \Psi^+| (|\beta|^2 + |\alpha|^2 e^{-i\Omega\tau}) e^{-i\Omega\tau} \cos \sqrt{2}g\tau \\ & + |\uparrow\uparrow\rangle_{AB} \langle \uparrow\uparrow| (|\beta|^2 \cos \sqrt{2}g\tau + |\alpha|^2 e^{-i\Omega\tau}) e^{-i\Omega\tau} \\ & - i(\alpha^* \beta |\downarrow\downarrow\rangle_{AB} \langle \Psi^+| + \alpha \beta^* |\Psi^+\rangle_{AB} \langle \downarrow\downarrow|) e^{-i\Omega\tau} \sin \sqrt{2}g\tau \\ & - i(\alpha \beta^* |\uparrow\uparrow\rangle_{AB} \langle \Psi^+| + \alpha^* \beta |\Psi^+\rangle_{AB} \langle \uparrow\uparrow|) e^{-i\Omega\tau} \sin \sqrt{2}g\tau, \end{aligned} \quad (12)$$

where $|\Psi^{\pm}\rangle_{AB} = (|\uparrow\downarrow\rangle_{AB} \pm |\downarrow\uparrow\rangle_{AB})/\sqrt{2}$ and $|\phi\rangle_X = \alpha|\uparrow\rangle_X + \beta|\downarrow\rangle_X$, with $|\uparrow\rangle_X(|\downarrow\rangle_X)$ the eigenstates of σ_3 belonging to the eigenvalues $+1$ (-1). Since the Hamiltonian (11) is symmetric under the exchange between A and B, $V_{\phi}(\tau)$ splits into the singlet and triplet sectors, and the singlet state $|\Psi^-\rangle_{AB}$ is one of the four eigenstates of $V_{\phi}(\tau)$. It is hence possible to extract a Bell state $|\Psi^-\rangle_{AB}$ provided (i) the corresponding eigenvalue

$$\lambda_{\Psi^-} = e^{-i\Omega\tau} (|\beta|^2 + |\alpha|^2 e^{-i\Omega\tau}) \quad (13)$$

is larger (in magnitude) than any other eigenvalue, and the extraction is efficient if the conditions (10), i.e. (ii) $|\lambda_{\Psi^-}| = 1$ and (iii) $|\lambda_n/\lambda_{\Psi^-}| \ll 1$ ($n \neq \Psi^-$), are fulfilled.

Condition (ii) is achieved by tuning τ such as

$$|\Omega|\tau = 2n\pi \quad (n = 0, 1, 2, \dots), \quad (14)$$

and condition (i) is possible unless $|\phi\rangle_X = |\uparrow\rangle_X$ or $|\downarrow\rangle_X$. Let us hence consider, under the condition (14), the case where

$$\alpha = \beta = 1/\sqrt{2}, \quad \text{i.e.} \quad |\phi\rangle_X = |\rightarrow\rangle_X \equiv (|\uparrow\rangle_X + |\downarrow\rangle_X)/\sqrt{2}. \quad (15)$$

In this case, three other eigenvalues than λ_{Ψ^-} are given by

$$\lambda_{\Phi^-} = \cos^2 \frac{g\tau}{\sqrt{2}}, \quad \lambda_{\pm} = 1 - \frac{1}{2} \sin \frac{g\tau}{\sqrt{2}} \left(3 \sin \frac{g\tau}{\sqrt{2}} \pm \sqrt{1 - 9 \cos^2 \frac{g\tau}{\sqrt{2}}} \right), \quad (16)$$

whose magnitudes are less than 1, fulfilling condition (i), when

$$|g|\tau/\sqrt{2} \neq m\pi/2 \quad (m = 0, 1, 2, \dots). \quad (17)$$

Therefore, the set of such parameters as (14), (15) and (17) enables us to extract the Bell state $|\Psi^-\rangle_{AB}$ from a general mixed state ϱ_{tot} .

The extraction of the Bell state $|\Psi^-\rangle_{AB}$ is demonstrated in figure 1, where the fidelity to the target state $|\Psi^-\rangle_{AB}$, defined by $F^{(\tau)}(N) \equiv {}_{AB} \langle \Psi^- | \varrho_{AB}^{(\tau)}(N) | \Psi^- \rangle_{AB}$, and




Figure 1. Extraction of a Bell state $|\Psi^-\rangle_{AB}$ from a product state $\varrho_{\text{tot}} = |\rightarrow\rangle_X \langle \rightarrow| \otimes |\uparrow\rangle_A \langle \uparrow| \otimes |\downarrow\rangle_B \langle \downarrow|$ [or from a mixed state $\varrho_{\text{tot}} = |\rightarrow\rangle_X \langle \rightarrow| \otimes (|\uparrow\downarrow\rangle_{AB} \langle \uparrow\downarrow| + |\downarrow\uparrow\rangle_{AB} \langle \downarrow\uparrow|)/2$]. Parameters are $\tau = 2\pi/\Omega$, $|\phi\rangle_X = |\rightarrow\rangle_X$ and $g = 0.25\Omega$ ($\Omega > 0$).

the success probability $P^{(\tau)}(N)$ versus the number of measurements, N , are shown for the initial state $\varrho_{\text{tot}} = |\rightarrow\rangle_X \langle \rightarrow| \otimes |\uparrow\rangle_A \langle \uparrow| \otimes |\downarrow\rangle_B \langle \downarrow|$, a (pure but) product state [or for a mixed state $\varrho_{\text{tot}} = |\rightarrow\rangle_X \langle \rightarrow| \otimes (|\uparrow\downarrow\rangle_{AB} \langle \uparrow\downarrow| + |\downarrow\uparrow\rangle_{AB} \langle \downarrow\uparrow|)/2$]. It is clear that the Bell state $|\Psi^-\rangle_{AB}$ is extracted after only 4 or 5 measurements (in this case, $|\lambda_{\Phi^-}| \simeq 0.20$ and $|\lambda_{\pm}| \simeq 0.44$). Since condition (ii) is fulfilled, the decay of the success probability $P^{(\tau)}(N)$ is suppressed, yielding $P^{(\tau)}(N) \rightarrow {}_{AB}\langle\Psi^-|\varrho_{AB}|\Psi^-\rangle_{AB} (= 1/2$ in figure 1), which means that the $|\Psi^-\rangle_{AB}$ component contained in the initial state ϱ_{AB} (after the zeroth measurement on ϱ_{tot}) is fully extracted. In this sense, the extraction is *optimal*.

4. Concluding remarks

The above example clearly and explicitly shows that our purification scheme works for entanglement purification. It is quite simple: one has simply to repeat one and the same measurement. At the same time, the purification can be made optimal: the optimal success probability is attainable as in the above example. Since the basic framework presented in section 2 is general, it possesses wide potential applicabilities in various settings for quantum computation and quantum information.

Acknowledgements

This work is partly supported by Grants-in-Aid for Scientific Research (C) from the Japan Society for the Promotion of Science (No. 14540280) and Priority Areas Research (B) from the Ministry of Education, Culture, Sports, Science and Technology, Japan (No. 13135221), by a Waseda University Grant for Special Research Projects (No. 2002A-567), and by the bilateral Italian-Japanese project 15C1 on ‘Quantum Information and Computation’ of the Italian Ministry for Foreign Affairs.

References

- [1] NIELSEN, M. A., and CHUANG, I. L., 2000, *Quantum Computation and Quantum Information* (Cambridge: Cambridge University Press), and references therein.
- [2] BOUWMEESTER, D., EKERT, A., and ZEILINGER, A., 2000, *The Physics of Quantum Information* (Heidelberg: Springer-Verlag), and references therein.
- [3] BENNETT, C. H., BRASSARD, G., POPESCU, S., SCHUMACHER, B., SMOLIN, J. A., and WOOTTERS, W. K., 1996, *Phys. Rev. Lett.*, **76**, 722; BENNETT, C. H., DIVINCENZO, D. P., SMOLIN, J. A., and WOOTTERS, W. K., 1996, *Phys. Rev. A*, **54**, 3824.
- [4] NAKAZATO, H., TAKAZAWA, T., and YUASA, K., 2003, *Phys. Rev. Lett.*, **90**, 060401.
- [5] For reviews, see: NAKAZATO, H., NAMIKI, M., and PASCAZIO, S., 1996, *Int. J. Mod. Phys. B*, **10**, 247; HOME, D., and WHITAKER, M. A. B., 1997, *Ann. Phys. (N.Y.)*, **258**, 237; FACCHI, P., and PASCAZIO, S., 2001, in *Progress in Optics*, edited by E. Wolf (Amsterdam: Elsevier), Vol. 42, p. 147.